



TRYVIO™

(aprocitentan) 12.5mg tablets



ARE YOU MEETING YOUR BLOOD PRESSURE GOALS?

Help lower your difficult-to-control hypertension by adding once-daily TRYVIO.

What is TRYVIO?

TRYVIO is a prescription medicine used to treat high blood pressure (hypertension) in adults who are taking other high blood pressure medicines and whose blood pressure is not well controlled. It is not known if TRYVIO is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TRYVIO?

TRYVIO may cause serious side effects, including:

Serious birth defects.

- TRYVIO may cause serious birth defects if taken during pregnancy.

Please see full Important Safety Information on pages 10 and 11. See full [Prescribing Information](#) and [Medication Guide](#).

If your blood pressure is still high, even with multiple medications, you're not alone.



3 out of **4** adults with high blood pressure still struggle to keep it under control, even when taking medications



Many people take **multiple medications**, yet their blood pressure is still not where it needs to be



Uncontrolled blood pressure **increases the risk** of heart attack, stroke, kidney disease and other serious health problems

IMPORTANT SAFETY INFORMATION

- Females who can become pregnant should not be pregnant when they start taking TRYVIO or become pregnant during treatment with TRYVIO or for 1 month after stopping treatment with TRYVIO.
- Females who can become pregnant should have a negative pregnancy test before starting treatment with TRYVIO.
- Females who **can become pregnant** are females who:
 - have entered puberty, even if they have not started their menstrual period, and
 - have a uterus, **and**
 - have not gone through menopause. Menopause means that you have not had a menstrual period for at least 12 months for natural reasons, or that you have had your ovaries removed.

Please see full Important Safety Information on pages 10 and 11. See full Prescribing Information and Medication Guide.

Why lowering blood pressure matters?



If you have been struggling with high blood pressure, the good news is—**even small improvements can help.**



Studies show that reducing blood pressure by just **5 mm Hg** may lower your chances of heart attack or stroke by about 10%.

That means every little step toward lowering your numbers really does count.


IMPORTANT SAFETY INFORMATION

- Females who cannot become pregnant are females who:
 - have not yet entered puberty, **or**
 - do not have a uterus, **or**
 - have gone through menopause. Menopause means that you have not had a menstrual period for at least 12 months for natural reasons, or that you have had your ovaries removed, **or**
 - are infertile for other medical reasons and this infertility is permanent and cannot be reversed.
- **Females who can become pregnant should use effective birth control during treatment with TRYVIO, and for 1 month after stopping TRYVIO** because the medicine may still be in your body.
 - Talk with your healthcare provider or gynecologist (a healthcare provider who specializes in female reproduction) to find out about options for effective birth control that you may use to prevent pregnancy during treatment with TRYVIO.
 - If you decide that you want to change the form of birth control that you use, talk with your healthcare provider or gynecologist to be sure that you choose another acceptable form of birth control.

TRYVIO helps take the pressure down


TRYVIO **works differently** than other blood pressure medications and may help lower your blood pressure when other treatments haven't been enough.

Hypertensive State



Overactive endothelin causes blood vessels to narrow and tighten, raising blood pressure

The TRYVIO Effect



TRYVIO targets endothelin and relaxes blood vessels to lower blood pressure

15.4
mm Hg*

In the clinical study[†], TRYVIO lowered systolic blood pressure by **15.4 mm Hg** after 4 weeks when added to other common blood pressure lowering medications.

Reduction in systolic blood pressure

* At week 4, significant reduction in sitting office SBP at week 4 by 15.4 mm Hg (97.5% CL, [-17.5, -13.3]) vs 11.6 mm Hg for placebo, for a difference of 3.8 mm Hg vs TRYVIO (97.5% CL, [-6.8, -0.8]; $P=0.0043$).

[†]TRYVIO was studied in a phase 3, multicenter trial involving 730 adults who had systolic blood pressure ≥ 140 mm Hg and were prescribed at least 3 antihypertensive medications which were continued throughout the study. The primary efficacy endpoint was the change in sitting systolic blood pressure from baseline to week 4, measured by unattended automated office blood pressure.

IMPORTANT SAFETY INFORMATION

- **Do not have unprotected sex.** Talk to your healthcare provider or pharmacist right away if you have unprotected sex or if you think your birth control has failed. Your healthcare provider may talk with you about using emergency birth control.
- **Tell your healthcare provider right away if you miss a menstrual period or you think you might be pregnant.**

Who should not take TRYVIO?

Do not take TRYVIO if you are:

- **Pregnant or currently trying to become pregnant.**
- **allergic to aprocitentan or any of the ingredients in TRYVIO.**

Please see full Important Safety Information on pages 10 and 11. See full [Prescribing Information](#) and [Medication Guide](#).

How to take TRYVIO



ONCE-DAILY
ONE 12.5-mg TABLET

Not actual size.

Once-daily TRYVIO is added on top of your current medications and can be taken:



With or without food



During day or night



With other medications

IMPORTANT SAFETY INFORMATION

Before taking TRYVIO, tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems
- have heart failure
- have anemia
- have kidney problems or get dialysis
- are pregnant or plan to become pregnant during treatment with TRYVIO. TRYVIO can cause serious birth defects.
- are breastfeeding or plan to breastfeed. It is not known if TRYVIO passes into your breastmilk. **Do not breastfeed** if you take TRYVIO. Talk to your healthcare provider about the best way to feed your baby if you take TRYVIO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

TRYVIO patient support



Insurance support

Help navigate your insurance benefits and support benefits verification



Financial support

Find the lowest out-of-pocket cost for your prescription

IMPORTANT SAFETY INFORMATION

What are the possible side effects of TRYVIO? TRYVIO may cause serious side effects, including:

- **Serious birth defects.**
- **Liver problems.** TRYVIO may cause liver problems. Your healthcare provider should do blood tests to check your liver before starting treatment and as needed during treatment with TRYVIO. Tell your healthcare provider if you have any of the following symptoms of liver problems during treatment with TRYVIO:
 - nausea or vomiting
 - yellowing of your skin or whites of your eyes
 - pain in the upper right stomach
 - dark urine
 - tiredness
 - fever
 - loss of appetite
 - itching

Please see full Important Safety Information on pages 10 and 11. See full [Prescribing Information](#) and [Medication Guide](#).

How to get TRYVIO

We've partnered with **Walgreens Specialty Pharmacy** to help make getting **TRYVIO simple and affordable.**

Your healthcare provider may send your prescription to Walgreens Specialty Pharmacy, which is different than your local retail store.

After your healthcare provider sends your TRYVIO prescription to Walgreens Specialty Pharmacy, you'll receive a phone call within 24 to 48 hours.

The call may come from a number you don't recognize—be sure to answer, so your prescription can be *processed without delay.*

During the call, they will:

1 Provide an overview of pharmacy services and support

2 Review insurance coverage and assess eligibility for savings programs

3 Confirm demographic and shipping information

To get started right away or if you haven't heard from Walgreens Specialty, call 888-347-3416

IMPORTANT SAFETY INFORMATION

- **Fluid retention.** Fluid retention and swelling are common during treatment with TRYVIO and can be serious. Tell your healthcare provider right away if you have any unusual weight gain, trouble breathing, or swelling of your ankles or legs. Your healthcare provider may treat you with other medicines (diuretics) if you develop fluid retention or swelling.
- **Low red blood cell levels (anemia).** Anemia is common during treatment with TRYVIO and can be serious. Your healthcare provider will do blood tests to check your red blood cells before starting and as needed during treatment with TRYVIO.



TRYVIO 30-day Free Trial

Get a 30-day free* trial of TRYVIO, regardless of insurance.

TRYVIO 30-DAY FREE TRIAL CARD

Group: AV44102003
BIN #: 019158
PCN: CNRX
ID: 69999613691

*Limitations apply. See full [Terms and Conditions](#).

FREE TRIAL TERMS AND CONDITIONS

This offer is not contingent on a purchase of any kind. The free trial may be redeemed for a single, one-time-only, 30-day supply (maximum 30 tablets) of TRYVIO. Patients must be new to TRYVIO and must not have previously filled a prescription for TRYVIO. A patient must be under the care of a licensed US physician. Patients must have a valid prescription(s) for TRYVIO. This offer may be used by cash-paying patients, patients with commercial insurance, and patients who are eligible for or participate in federal healthcare programs such as Medicaid, Medicare, or any similar federal or state programs only when patients, pharmacists, and prescribers agree not to seek reimbursement from health insurance, health savings or flexible spending accounts, or any third party, including state or federally funded programs for the free trial of TRYVIO received by the patient through this offer. Patients eligible for the free trial may not count the free trial as an expense incurred for determining out-of-pocket costs for any plan, including true out-of-pocket costs ("TrOOP"), under Medicare Part D. This offer is limited to one per patient and is nontransferable. This free trial offer cannot be combined with any other free trial, coupon, discount, prescription savings card or other offer. No substitutions are permitted. This offer is not health insurance, nor is it provided for the purpose of financial assistance. This offer is restricted to residents of the United States. Void where prohibited by law, taxed or restricted. Not valid in California or Massachusetts, if an AB-rated generic equivalent becomes available for the product. This offer may be changed or discontinued at any time without notice. Patients are responsible for applicable taxes, if any. By redeeming this free trial voucher, you acknowledge that you are an eligible patient and that you understand and agree to comply with the terms and conditions of this offer.

For Healthcare Professionals Only: For any questions regarding SS&C online processing, please call the Help Desk at **844-373-0987**.

IMPORTANT SAFETY INFORMATION

- **Decreased sperm count.** TRYVIO may cause decreased sperm counts in males and may affect the ability to father a child. Tell your healthcare provider if being able to have children is important to you.

Please see full Important Safety Information on pages 10 and 11. See full [Prescribing Information](#) and [Medication Guide](#).

TRYVIO Copay Savings Program

Commercially insured eligible patients may receive TRYVIO for as little as \$10* per month.



TRYVIO \$10 SAVINGS CARD

Group: EC44102002

BIN #: 019158

PCN: CNRX

ID: 09998483288

*Limitations apply. See full [Terms and Conditions](#).

TRYVIO SAVINGS CARD TERMS OF USE AND RESTRICTIONS

Eligibility Requirements: Patients may be eligible if: (1) the patient is insured by commercial insurance and their prescription insurance coverage does not cover the full cost of the prescription; (2) patient does not have prescription insurance coverages through a state or federal healthcare program, including but not limited to Medicare Part D, Medicaid, Medigap, Veterans Affairs (VA), or Department of Defense (DOD) programs; patients who move from commercial insurance plans to state or federal healthcare programs will no longer be eligible; (3) patient is 18 years of age or older; (4) patient is a resident of the United States.

Terms of Use: Eligible commercially insured patients with a valid prescription for TRYVIO (aprocitentan) 12.5 mg who present this savings card at participating pharmacies may pay as little as \$10 for a 30-day supply. Maximum savings limit applies; patient out-of-pocket expenses may vary. The patient is responsible for applicable taxes—void where prohibited by law, taxed, or restricted. Other restrictions may apply. Patients, pharmacists, and prescribers cannot seek reimbursement from health insurance or any third party for any part of the benefit received by the patient through this offer. All copay payments are for the benefit of the patient only. Idorsia reserves the right to rescind, revoke, or amend this offer, eligibility, and terms of use at any time without notice. Must present offer along with a valid prescription at the time of purchase.

Restrictions: This offer is not valid for cash-paying patients. This offer is nontransferable, no substitutions are permissible, and the offer cannot be applied with any other financial assistance program, free trial, discount, prescription savings card, or other offers. The savings card for TRYVIO is not health insurance. The savings card may not be sold, purchased, or traded. ConnectiveRx manages this program on behalf of Idorsia.

For Healthcare Professionals Only: Pharmacist Instructions for a Patient with an Eligible Third Party: For Insured/Covered Patients: Submit the claim to the primary Third-Party Payer first, then submit the balance due to SS&C as a Secondary Payer COB with patient responsibility amount and a valid Other Coverage Code of 08. Patient pays \$10 for a 30-day supply. The pharmacist will receive reimbursement from SS&C. For any questions regarding SS&C online processing, please call the Help Desk at 844-373-0987.

IMPORTANT SAFETY INFORMATION

Your healthcare provider may stop treatment with TRYVIO if you develop certain side effects. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of TRYVIO. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Indication and Important Safety Information

What is TRYVIO?

TRYVIO is a prescription medicine used to treat high blood pressure (hypertension) in adults who are taking other high blood pressure medicines and whose blood pressure is not well controlled. It is not known if TRYVIO is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TRYVIO?

TRYVIO may cause serious side effects, including:

Serious birth defects.

- **TRYVIO may cause serious birth defects if taken during pregnancy.**
- **Females who can become pregnant should not be pregnant when they start taking TRYVIO or become pregnant during treatment with TRYVIO or for 1 month after stopping treatment with TRYVIO.**
- **Females who can become pregnant should have a negative pregnancy test** before starting treatment with TRYVIO.
- Females who **can become pregnant** are females who:
 - have entered puberty, even if they have not started their menstrual period, and
 - have a uterus, **and**
 - have not gone through menopause. Menopause means that you have not had a menstrual period for at least 12 months for natural reasons, or that you have had your ovaries removed.
- Females who cannot become pregnant are females who:
 - have not yet entered puberty, **or**
 - do not have a uterus, **or**
 - have gone through menopause. Menopause means that you have not had a menstrual period for at least 12 months for natural reasons, or that you have had your ovaries removed, **or**
 - are infertile for other medical reasons and this infertility is permanent and cannot be reversed.
- **Females who can become pregnant should use effective birth control during treatment with TRYVIO, and for 1 month after stopping TRYVIO** because the medicine may still be in your body.
 - Talk with your healthcare provider or gynecologist (a healthcare provider who specializes in female reproduction) to find out about options for effective birth control that you may use to prevent pregnancy during treatment with TRYVIO.
 - If you decide that you want to change the form of birth control that you use, talk with your healthcare provider or gynecologist to be sure that you choose another acceptable form of birth control.
- **Do not have unprotected sex.** Talk to your healthcare provider or pharmacist right away if you have unprotected sex or if you think your birth control has failed. Your healthcare provider may talk with you about using emergency birth control.
- **Tell your healthcare provider right away if you miss a menstrual period or you think you might be pregnant.**

Who should not take TRYVIO?

Do not take TRYVIO if you are:

- **Pregnant or currently trying to become pregnant.**
- **allergic to aprocitentan or any of the ingredients in TRYVIO.**

Before taking TRYVIO, tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems
- have heart failure
- have anemia
- have kidney problems or get dialysis
- are pregnant or plan to become pregnant during treatment with TRYVIO. TRYVIO can cause serious birth defects.
- are breastfeeding or plan to breastfeed. It is not known if TRYVIO passes into your breastmilk. **Do not breastfeed** if you take TRYVIO. Talk to your healthcare provider about the best way to feed your baby if you take TRYVIO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of TRYVIO? TRYVIO may cause serious side effects, including:

- **Serious birth defects.**
- **Liver problems.** TRYVIO may cause liver problems. Your healthcare provider should do blood tests to check your liver before starting treatment and as needed during treatment with TRYVIO. Tell your healthcare provider if you have any of the following symptoms of liver problems during treatment with TRYVIO:
 - nausea or vomiting
 - yellowing of your skin or whites of your eyes
 - pain in the upper right stomach
 - dark urine
 - tiredness
 - fever
 - loss of appetite
 - itching
- **Fluid retention.** Fluid retention and swelling are common during treatment with TRYVIO and can be serious. Tell your healthcare provider right away if you have any unusual weight gain, trouble breathing, or swelling of your ankles or legs. Your healthcare provider may treat you with other medicines (diuretics) if you develop fluid retention or swelling.
- **Low red blood cell levels (anemia).** Anemia is common during treatment with TRYVIO and can be serious. Your healthcare provider will do blood tests to check your red blood cells before starting and as needed during treatment with TRYVIO.
- **Decreased sperm count.** TRYVIO may cause decreased sperm counts in males and may affect the ability to father a child. Tell your healthcare provider if being able to have children is important to you.

Your healthcare provider may stop treatment with TRYVIO if you develop certain side effects. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of TRYVIO. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please read the [Medication Guide](#) for TRYVIO and discuss it with your healthcare provider.

Please see full Prescribing Information and Medication Guide.

Highlights of TRYVIO

Help lower difficult-to-control hypertension by adding TRYVIO



1 tablet, taken once-daily



Helps reduce high blood pressure



Can be taken with other medications



Learn more about TRYVIO
by visiting
www.tryvio.com

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TRYVIO?

TRYVIO may cause serious side effects, including:

Serious birth defects.

- TRYVIO may cause serious birth defects if taken during pregnancy.
- Females who can become pregnant should not be pregnant when they start taking TRYVIO or become pregnant during treatment with TRYVIO or for 1 month after stopping treatment with TRYVIO.
- Females who can become pregnant should have a negative pregnancy test before starting treatment with TRYVIO.

Please see full Prescribing Information and Medication Guide.